


Alkynes

Alkynes or acetylenes are compounds that contain a carbon-carbon triple bond.

E.g.



acetylene

The triple bond results in a molecular formula of $\text{C}_n\text{H}_{2n-2}$


Ethane	C_2H_6	0 elements of unsaturation
Ethene	C_2H_4	1 element of unsaturation
Ethyne	C_2H_2	2 elements of unsaturation

The triple bond contributes two elements of unsaturation.


Nomenclature of Alkynes

IUPAC nomenclature is similar to that for alkenes, except the -ane ending is replaced with -yne.

The chain is numbered from the end closest to the triple bond.




When additional functional groups are present, the suffixes are combined.




Terminal and Internal Alkynes

The position of the triple bond can alter the reactivity of the alkyne.

Compounds with triple bonds at the end of a molecule are called terminal alkynes. (Terminal C-H groups are called acetylenic hydrogens).



terminal
alkyne



internal
alkyne

If the triple bond is flanked by alkyl groups on both sides it is an internal alkyne.

Electronic Structure of Alkynes

The triple bond consists of one σ bond and two π bonds.


Each carbon is bonded to two other atoms, and there are no nonbonding electrons.

Carbon requires two hybrid orbitals to bond to the atoms, thus sp hybrids are used.

The sp orbitals are linear and oriented at 180° .


The C-C bond is formed from sp - sp overlap

The C-H bond is formed from sp - s overlap.



The creation of sp hybrids leaves 2 free p orbitals – these create the two π bonds.

The p orbitals overlap just as for an alkene, except there are two π bonds at right angles to one another.




This generates a cylinder of electron density around the sigma bond.

The C-C bond length for ethyne is 1.20\AA which is shorter than ethane (1.54\AA) and ethene (1.33\AA).

The C-H bond length in ethyne is 1.06\AA which is also shorter than in ethane (1.09\AA) or ethene (1.08\AA).




This is because the C-H bond contains more s character ($sp^3 \rightarrow sp^2 \rightarrow sp$) which gives stronger bonds.



Acidity of Alkynes



Terminal alkynes are acidic, the end hydrogen can be removed as a proton by a strong base.

E.g.




Heavy Metal Acetylides

These form more covalent type structures, and the silver and copper acetylides will precipitate from solution. (Test for terminal alkynes).




Synthesis of Alkynes

The carbanions produced are called acetylides, and are strong nucleophiles.




They can react in $\text{S}_{\text{N}}2$ displacements of halides or tosylates from primary unhindered substrates to produce internal alkynes.

E.g.




Carbonyl and Epoxide Reactions




Acetylides will also attack other standard electrophiles like carbonyls and epoxides to generate alcohols. (These reactions are very similar to Grignard type reactions).

E.g.




Epoxides also generate longer chain alcohols.




Synthesis of Alkynes via Elimination

In the same way that alkenes are produced by elimination of H-X, alkynes can be produced by elimination of 2 moles of H-X from a geminal or vicinal dihalide.



Strongly basic conditions (KOH or NaNH₂, high temps) are required for this transformation, and often the molecule may not be stable to these conditions and may be destroyed or will rearrange.


Base Catalyzed Rearrangements



Often the deprotonated acetylide will rearrange to give a more stable product.


More highly substituted alkynes are more stable.

Sodium amide is such a strong base that during the equilibrating of all triple bond isomers, the terminal alkyne gets deprotonated and exists as the sodium salt, and is therefore prevented from equilibrating further.



Eventually all the alkynes are trapped as the terminal sodium acetylide salts, and on addition of water generates the terminal alkyne.

E.g.




Bond Strengths

Recall that the π bond of an alkene is weaker than its σ bond.

Compound	Total Energy	Class	Approx. Energy
Alkane	83kcal/mol	alkane σ	83kcal/mol
Alkene	146kcal/mol	alkene π	63kcal/mol
Alkyne	200kcal/mol	alkyne 2 nd π	54kcal/mol

The sum of the two π bonds in an alkyne are **weaker** than the sum of two alkene π bonds.

The triple bond is **electron rich** and acts as a nucleophile (just like a double bond).




Since addition reactions convert π bonds into σ bonds, these reactions are typically exothermic.

Alkynes can add up to two molecules across the triple bond.

Addition of Hydrogen


Hydrogen in the presence of a catalyst will add twice to alkynes to generate alkanes.



This reaction proceeds through a cis alkene intermediate, but **cannot** be stopped at this stage except with the use of a special catalyst.

Lindlar's catalyst


This is a partially deactivated (poisoned) catalyst consisting of barium sulfate, palladium and quinoline (the poison).




The hydrogens are delivered simultaneously to the same side of the alkyne, creating syn addition (cis alkenes).

Trans Alkenes

These can be produced by the action of sodium metal in liquid ammonia.



An electron is added to the triple bond, generating a radical anion, which is protonated to generate a vinyl radical.




The radical accepts another electron which produces an anion, which in turn gets protonated, giving the alkene with **anti** stereochemistry.

The anti stereochemistry arises from the two alkyl groups being as far away from each other as possible at the vinyl radical stage.


Addition of Halogens

Chlorine and bromine add in an analogous fashion as for alkenes.

If only one mole of halogen is added then a dihaloalkene is produced, and a mixture of syn and anti addition is observed.




It is usually hard to control the addition of just one equivalent of halogen and it is more common to add two equivalents to generate tetrahalides.




Addition of Hydrogen Halides

H-X can be added to alkynes just like to alkenes.




For **terminal** alkynes, the products have orientation as dictated by Markovnikov's rule.

The vinyl cation is more stable with positive charge on the **more** substituted carbon.




The second mole of H-X will usually add with the same orientation as the first.

Internal alkynes show little selectivity since both intermediate cations are equally substituted.



The peroxide effect is also observed with the addition of H-Br and peroxides to generate anti-Markovnikov products.




Hydration of Alkynes to Ketones and Aldehydes

There are two different methods:

Mercuric-ion catalyzed hydrations


A mixture of mercuric sulfate and aqueous sulfuric acid is used to perform addition of water across the triple bond.

The orientation is Markovnikov, and the initial products are enols.



Electrophilic addition of the Hg^{2+} ion generates a vinyl cation, which is attacked by a nucleophilic water molecule.


Deprotonation gives an organomercurial alcohol which, under acidic conditions, has the mercury replaced by a hydrogen, to generate a vinyl alcohol (enol).




Enols tend to be unstable, and isomerise to their more stable ketone form.

Oxidation of Alkynes


Under mild (cold, neutral) conditions the reaction of potassium permanganate with an alkyne can generate a diketone.



If the mixture becomes too warm or basic, the oxidation proceeds further to generate two carboxylate anions, which on acidification generates two carboxylic acids.



Terminal alkynes are similarly cleaved to give a carboxylic acid and carbon dioxide.



Ozonolysis

Ozonolysis of an alkyne followed by hydrolysis gives similar products to those obtained from permanganate oxidative cleavage.

